

People who received COVID-19 vaccine outside the United States - Quick Reference

Vaccination Status	Primary Series	Immunocompromised Individuals	Boosters
People who completed all the recommended doses of a *WHO-EUL COVID-19 vaccine or people who completed a heterologous (mix and match) series composed of any combination of FDA-approved, FDA-authorized, or *WHO-EUL COVID-19 vaccines	Individuals with normal immune systems are fully vaccinated and no further primary dose needed	Immunocompromised Individuals are fully vaccinated if: Individuals aged ≥ 12 years receive an additional primary dose of Pfizer COVID-19 Vaccine (30 μg) at least 28 days after receiving the second vaccine dose	Individuals aged ≥ 16 years (including immunocompromised people who received an additional primary dose) should receive a single booster dose of Pfizer COVID-19 Vaccine (30 μg) at least 6 months after completing their primary series.
People who received only the first dose of a multidose *WHO-EUL COVID-19 primary series that is not FDA-approved or FDA-authorized, or received COVID-19 vaccine primary series that is not listed for emergency use by *WHO	Individuals should restart primary series with an FDA-approved or authorized Covid-19 Vaccine, with a minimum interval of at least 28 days since receipt of the last dose of a non-FDA approved/authorized vaccine	Additional dose is NOT recommended at this time. <i>After completion of primary vaccination with an FDA-approved or FDA-authorized COVID-19 vaccine, these individuals are considered fully vaccinated, and are not recommended to receive an additional primary dose at this time.</i>	Booster dose is NOT recommended at this time.

*WHO COVID-19 vaccines for emergency use include:

- Pfizer-BioNTech COVID-19 Vaccine (e.g., BNT162b2, COMIRNATY, Tozinameran)
- Janssen (Johnson & Johnson) COVID-19 Vaccine (e.g., Ad26.COV2.S)
- Moderna COVID-19 Vaccine (e.g., mRNA 1273, Takeda, Spikevax)
- Sinopharm-BIBP COVID-19 Vaccine
- Sinovac-CoronaVac COVID-19 Vaccine
- Bharat Biotech International COVID-19 Vaccine (e.g., BBV152, COVAXIN)
- AstraZeneca-Oxford COVID-19 Vaccine (e.g., [ChAdOx1-S (recombinant)], AZD1222, Covishield, Vaxzevria)

Individuals who were vaccinated outside the United States with a currently FDA-approved or FDA-authorized COVID-19 vaccine (Pfizer, Moderna, or J&J), should follow the same protocols as if they were vaccinated in the United States. See page 2

Individuals who were vaccinated outside the United States with a currently FDA-approved or FDA-authorized COVID-19 vaccine (Pfizer, Moderna, or J&J), should follow the same protocols as if they were vaccinated in the United States.

Vaccination Status	Primary Series	Immunocompromised Individuals	Boosters
<p>Individuals who were vaccinated outside the United States with a currently FDA-approved or FDA-authorized COVID-19 vaccine (Pfizer, Moderna, or J&J). ++No changes to recommended</p>	<p>Individuals with normal immune systems are fully vaccinated and no further primary dose is needed if they received:</p> <ul style="list-style-type: none"> • single dose J&J or • 2 doses of Pfizer 30 µg IM or Moderna 100 µg IM <p>If the individual only received 1 dose of Pfizer or Moderna, provide 2nd dose (do not restart series)</p>	<p>Immunocompromised Individuals are fully vaccinated if:</p> <p>Individuals aged ≥12 years receive an additional dose of Pfizer COVID-19 Vaccine (30 µg) at least 28 days after receiving the second vaccine dose.</p> <p>Individuals aged ≥18 years receive an additional dose of Moderna COVID-19 Vaccine (100 µg) at least 28 days after receiving the second vaccine dose.</p> <p>A single dose of J&J.</p>	<p>Individuals aged ≥16 years (including immunocompromised people who received an additional primary dose) may receive a booster, at least 6 months after completing their primary series of Pfizer or Moderna’s COVID-19 vaccine, or at least two months after the J&J COVID-19 Vaccine.</p> <p>Individuals may choose which vaccine they receive as a booster dose.</p> <p>Pfizer-BioNTech = 16 years of age or older = 30 µg IM Moderna = 18 years of age or older = 50 µg IM (half dose)</p>